

Human Pulmonary Lymphatic Microvascular Endothelial Cells

Catalog Number	ALHE45
Product Name	Human Pulmonary Lymphatic Microvascular Endothelial cells
Storage	Liquid Nitrogen Vapor phase
Product Format	Frozen vial
Cells Number	500.000 cells or >90% confluent in frozen vial

***Caution:** The handling of human derived products has the potential to be biologically hazardous. All Cell strains tested negative for HIV, HBV, and HCV DNA in diagnostic tests. Proper precautions must be taken to avoid exposure. Always wear proper protective equipment (Gloves, safety glasses, etc.) when handling these materials. We recommend following the universal procedures for handling products of human origin as the minimum precaution against contamination.

GENERAL INFORMATION

HPuLyMVECS Cat.#ALHE45 cells are derived from healthy adult human lung tissue. The cells are shipped in frozen vials (the cells are provided @ passage 3). Endothelial Growth Medium (EGM, contains 10% serum and growth supplements, Cat.#EGM-2102) is recommended for cell culture and these cells have a minimum average population doubling levels > 10 when cultured following the detailed protocol described below).

CELL LINE CHARACTERIZATION

Cytoplasmic LYVE-1 and Podoplanin	>95% positive by immunofluorescence
Neuclis PROX-1	>95% positive by immunofluorescence
Cytoplasmic VWF / Factor VIII:	>95% positive by immunofluorescence
Cytoplasmic uptake of Di-I-Ac-LDL:	>95% positive by immunofluorescence
Cytoplasmic Cytoplasmic PECAM1	>95% positive by immunofluorescence
HPuLyMVECs are negative for	HIV-1, HBV, HCV, and mycoplasma

PRODUCT USE AND SHIPPING STATUS

Product Use	HPuMVECs are for research use only
Shipping Status	Frozen vial

Frozen Cell Line Sub Culture:

- 1) Coating T25 flasks. Add 2 ml AlphaBioCoat (Cat.#AC001) into a T25 flask and ensure entire interior surface is coated with solution. After 30 minutes, dispose of AlphaBioCoat by aspiration. Gently rinse and aspirate flask with Phosphate Buffer Solution (Cat.#1XPBS-001). The flask is now ready for use (no need for overnight incubation when coated with Cat.#AC001).
- 2) If you are using the coated flask the same day, add about 4 ml of Endo-Growth media (Cat.#EGM-2102) to the coated flask. If the media changes color from pink to yellow, aspirate and discard the media. Add 4ml of fresh media to the coated flask.
- 3) Thaw the cells in a 37°C water bath. Once you see a small amount of ice left in the vial, spray the vial with 70% Ethanol and wipe it down.
- 4) Transfer the vial into your Biosafety cabinet.
- 5) Using a 2 or 5ml pipet, pipet the cells out of the vial.
- 6) Transfer your cell suspension in to your coated plate that have the 4 ml media in it.
- 7) You should have a total working volume of 5ml of cell suspension in the flask; close the cap. Make sure cells are evenly distributed in the flask by moving the flask left and right five times. Move it up and down for and additional five times.
- 8) Place flask in a 37°C incubator with 5% CO₂. If flask is not vented, please loosen cap.
- 9) Change media after 48 hours.
- 10) For assay or to induce quiescent cell, use our Endothelial Basal Medium Cat.#EDBM-2101